



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

If  $\cos\phi = \infty$ ,  $\tan\phi = \pm i$ .

If  $\sin 15\phi = 0$ ,  $\phi = \frac{n\pi}{15}$  ( $n=0, 1, 2, \dots, 14$ ).

$\therefore x = \pm i, 0, \tan \frac{\pi}{15}, \tan \frac{2\pi}{15}, \dots, \tan \frac{14\pi}{15}$ ,

$$\begin{aligned}
 &= \pm i, 0, \pm \frac{\sqrt{[10+2\sqrt{5}]} - \sqrt{(15)} + \sqrt{(3)}}{\sqrt{[30+6\sqrt{5}]} + \sqrt{(5)} - 1}, \pm \frac{\sqrt{(15)} + \sqrt{(3)} - \sqrt{[10-2\sqrt{5}]}}{\sqrt{[30-6\sqrt{5}]} + \sqrt{(5)} + 1}, \\
 &\pm \frac{\sqrt{[10-2\sqrt{5}]}}{\sqrt{(5)} + 1}, \pm \frac{\sqrt{[10+2\sqrt{5}]} + \sqrt{(15)} - \sqrt{(3)}}{\sqrt{[30+6\sqrt{5}]} - \sqrt{(5)} + 1}, \pm \sqrt{(3)}, \\
 &\pm \frac{\sqrt{[10+2\sqrt{5}]}}{\sqrt{(5)} - 1}, \pm \frac{\sqrt{(15)} + \sqrt{(3)} + \sqrt{[10-2\sqrt{5}]}}{\sqrt{[30-6\sqrt{5}]} - \sqrt{(5)} - 1}.
 \end{aligned}$$

(These values are taken from Hobson's *Plane Trigonometry*.)

Professor J. Scheffer gave a partial solution, following the method in the above solution. Mr. Spuner sent in a similar solution, with results given in decimals instead of radicals.

**347. Proposed by GUSTAVE JACOBSON, A. M., Public Accountant, Chicago, Ill.**

A corporation needing some additional capital for a short term of years, issues \$300,000 of debenture bonds carrying 6% interest, and payable 1/5 each year for 5 years. Coupons are attached to the bonds maturing every six months; the bonds are sold at 90 flat. What average rate of interest does the company pay for the money, including interest on interest?

**Solution by THEODORE L. DeLAND, Treasury Department, Washington, D. C.**

Let  $2x$  = the rate of interest paid per annum, payable semi-annually; then  $x$  = the rate of semi-annual interest, compound. Only \$270,000 was used by the corporation, which was repaid, interest in 10 payments and principal in 5 payments; and they alternate, the first each six months and the second at the end of each year as follows: (1), \$9,000; (2), \$69,000, (3), \$7,200; (4), \$67,200; (5), \$5,400; (6), \$65,400; (7), \$3,600; (8), \$63,600; (9), \$1,800, and (10), \$61,800.

The problem may now be treated, algebraically, as a question in partial payments under the U. S. Court Rule; and we have, after taking out the factor 600, arranging, dropping the dollar sign as follows:

$$\begin{aligned}
 450(1+x)^{10} - [15(1+x)^9 + 115(1+x)^8 + 12(1+x)^7 + 112(1+x)^6 \\
 + 9(1+x)^5 + 109(1+x)^4 + 6(1+x)^3 + 106(1+x)^2 + 3(1+x)] = 103.
 \end{aligned}$$

Observe that as arranged the first member has 2 terms, and their difference

=103, As the debenture was discounted, the rate realized on the investment must be greater than 3% as a semi-annual rate. We try 5% and find it too small; we then try 5.0275%, with  $(1+x)=1.050275$ , and find the first member=102.9337, which is too small; we then try  $(1+x)=1.050285$  and obtain the value 102.9531, still too small. By double position we have a corrected value and  $(1+x)=1.050309$  and obtain 103.0304, which is now too large. By double position we have  $(1+x)=1.050291$ , and obtain 102.9937, again too small. By double position again we have  $(1+x)=1.0502941$  and obtain 103.0033, now too large. We now again use double position and obtain,  $(1+x)=1.05029269$ , this will give a value which will differ by less than 1% of a mill. We now have for a good approximation a semi-annual rate, 5.029269%; or the rate per annum payable semi-annually, 10.058538%.

Also solved by S. A. Corey who got for a result 10.28 per cent.; A. H. Holmes, whose result was 10.07 per cent., and the proposer, whose result was 10.0532 per cent. These contributors used practically the same method of solution. Mr. Spuner also sent in an excellent solution of 345, which reached us too late for credit in the January number.

## CALCULUS.

298. Proposed by C. N. SCHMALL, New York City.

Prove, by calculus, that if two regular polygons have equal perimeters, that which has the greater number of sides has the greater area.

Solution by E. L. SHERWOOD, Shady Side Academy, Pittsburg, Penn., and the PROPOSER.

In any regular polygon, let  $2p$ =the perimeter,  $n$ =number of sides.

$\therefore \frac{2p}{n}$ =each side, and  $\frac{p}{n \tan(\pi/n)}$ =the apothem.

$$\frac{p^2}{n \tan(\pi/n)} = \frac{p^2}{n} \cot \frac{\pi}{n} = \text{area} \dots (1).$$

Put  $u = n \tan \frac{\pi}{n} \dots (2)$ ; hence the area will be a maximum when  $u$  is a minimum.

From (2) we have

$$\frac{du}{dn} = \tan \frac{\pi}{n} - \frac{\pi}{n} \sec^2 \frac{\pi}{n} = \tan \phi - \phi \sec^2 \phi \quad (\text{where } \frac{\pi}{n} = \phi) = \sin \phi \sec \phi - \phi \sec^2 \phi$$

$$= \frac{1}{2} (2 \sin \phi \sec \phi - 2 \phi \sec^2 \phi) = \frac{1}{2} \sec^2 \phi \left( \frac{2 \sin \phi}{\sec \phi} - 2 \phi \right) = \frac{1}{2} \sec^2 \phi (\sin 2\phi - 2\phi),$$

$$= \frac{1}{2} \sec^2 \frac{\pi}{n} \left( \sin \frac{2\pi}{n} - \frac{2\pi}{n} \right).$$